

What does a safe sleep environment look like?

Reduce the risk of Sudden Infant Death Syndrome (SIDS) and other sleep-related causes of infant death



Use a firm sleep surface, such as a mattress in a safety-approved* crib, covered by a fitted sheet.

Do not use pillows, blankets, sheepskins, or crib bumpers anywhere in your baby's sleep area.

Keep soft objects, toys, and loose bedding out of your baby's sleep area.

Do not smoke or let anyone smoke around your baby.

Make sure nothing covers the baby's head.

Always place your baby on his or her back to sleep, for naps and at night.

Dress your baby in light sleep clothing, such as a one-piece sleeper, and do not use a blanket.

Baby should not sleep in an adult bed, on a couch, or on a chair alone, with you, or with anyone else.

*For more information on crib safety guidelines, contact the Consumer Product Safety Commission at 1-800-638-2772 or www.cpsc.gov



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
NATIONAL INSTITUTES OF HEALTH
Eunice Kennedy Shriver National Institute of Child
Health and Human Development



Eunice Kennedy Shriver
NICHD
National Institute of Child Health
& Human Development

Safe sleep for your baby



- Always place your baby on his or her back to sleep, for naps and at night, to reduce the risk of SIDS.
- Use a firm sleep surface, covered by a fitted sheet, to reduce the risk of SIDS and other sleep-related causes of infant death.
- Your baby should not sleep in an adult bed, on a couch, or on a chair alone, with you, or with anyone else.
- Keep soft objects, toys, and loose bedding out of your baby's sleep area to reduce the risk of SIDS and other sleep-related causes of infant death.
- To reduce the risk of SIDS, women should:
 - Get regular healthcare during pregnancy, and
 - Not smoke, drink alcohol, or use illegal drugs during pregnancy or after the baby is born.
- To reduce the risk of SIDS, do not smoke during pregnancy, and do not smoke or allow smoking around your baby.
- Breastfeed your baby to reduce the risk of SIDS.
- Give your baby a dry pacifier that is not attached to a string for naps and at night to reduce the risk of SIDS. Wait until your baby is breastfeeding well before trying a pacifier.
- Do not let your baby get too hot during sleep.
- Follow healthcare provider guidance on your baby's vaccines and regular health checkups.
- Avoid products that claim to reduce the risk of SIDS and other sleep-related causes of infant death.
- Do not use home heart or breathing monitors to reduce the risk of SIDS.
- Give your baby plenty of Tummy Time when he or she is awake and when someone is watching.



Remember Tummy Time!

Place babies on their stomachs when they are awake and when someone is watching. Tummy Time helps your baby's head, neck, and shoulder muscles get stronger and helps to prevent flat spots on the head.



To learn more or get help making a safe infant sleep area in your home:
HEALTH Information Line: 401-222-5960 / RI Relay 711
www.health.ri.gov/safesleep